

Calculus 1 — Exam 3
MAT 250, Spring 2012 — D. Ivanšić

Name: _____
Show all your work!

Differentiate and simplify where appropriate:

1. (6pts) $\frac{d}{du} (e^{2u} \cos u) =$

2. (4pts) $\frac{d}{dx} e^{\tan x} =$

3. (7pts) $\frac{d}{d\theta} (\sec^2 \theta - \tan^2 \theta) =$

4. (7pts) $\frac{d}{dx} \cos(\sqrt[3]{x^3 - 3x^2 + 14}) =$

5. (8pts) $\frac{d}{d\theta} \frac{\sin \theta}{\cos^2 \theta} =$

6. (8pts) Use implicit differentiation to find y' .

$$\sin(x + y) = x \cos y$$

7. (10pts) The circle of radius 3 centered at the origin has equation $x^2 + y^2 = 9$. Use implicit differentiation to find the equation of the tangent line at point $(-\sqrt{3}, -\sqrt{6})$. Draw the picture of the circle and the tangent line.

8. (14pts) A lemon is thrown from ground level at initial velocity 20m/s.
- Write the formula for the position of the lemon at time t (you may assume $g \approx 10$).
 - Write the formula for the velocity of the lemon at time t .
 - What is the lemon's velocity when it is at its highest point?
 - When does the lemon reach its highest point? What is the highest point?
 - What is the acceleration when the lemon is at its highest point? How about at time $t = 0$?

9. (14pts) If you bungee-jump with a cord of length x meters, the time in seconds you will spend free-falling (that is, until the bungee cord engages) is given by $t = \frac{\sqrt{x}}{\sqrt{5}}$.

- a) Find the free-falling time for a bungee cord of length 45m.
- b) Find the ROC of time with respect to cord length when $x = 45$ (units?).
- c) Use b) to estimate the change in time if cord length increases by 10m.
- d) Use c) to estimate the free-falling time for $x = 55$ m and compare to the actual value of 3.3166s.

10. (12pts) Let $f(x) = x^{-4}$.

- a) Find the first four derivatives of f .
- b) Find the general formula for $f^{(n)}(x)$.

11. (10pts) A remote-controlled bunny's position $s(t)$ is tracked for 10 seconds. Draw the graph of its position function if we know the following:

- $s(0) = 0$, $s(10) = 4$
- it moves forward on interval $(0, 6)$
- it moves backwards on interval $(6, 10)$
- it accelerates on interval $(0, 3)$
- it decelerates on interval $(3, 6)$
- it moves at a steady velocity on interval $(6, 10)$.

Bonus. (10pts) Let $f(x) = x^2e^x$.

- a) Find the first five derivatives of f .
- b) Find the pattern for $f^{(n)}(x)$.