

Final answers should have accuracy to 6 decimal places (or 4 decimal places for table-derived answers). Show some work how the mean and standard deviation are computed. *Giving only the answer will bring you few points.*

$$\begin{aligned} \text{midrange} &= \frac{\text{lowest value} + \text{highest value}}{2} & \text{range} &= \text{highest value} - \text{lowest value} \\ \bar{x} &= \frac{x_1 + x_2 + \cdots + x_n}{n} = \frac{\sum_i x_i}{n} = \frac{\sum_i f_i x_i}{n} & Z &= \frac{X - \bar{x}}{s} \\ s &= \sqrt{\frac{(x_1 - \bar{x})^2 + (x_2 - \bar{x})^2 + \cdots + (x_n - \bar{x})^2}{n - 1}} = \sqrt{\frac{\sum_i (x_i - \bar{x})^2}{n - 1}} = \sqrt{\frac{\sum_i f_i (x_i - \bar{x})^2}{n - 1}} \end{aligned}$$

1. (18pts) Over the course of two weeks (workdays only) a first-grader counts the number of toys left on the floor after recess. She gets the numbers below.

a) Find the midrange.

b) Find the median.

3, 2, 6, 4, 5, 5, 3, 1, 3, 4

c) Find the mean.

d) Find the range.

e) Find the standard deviation.

2. (8pts) A city is considering whether to renovate and expand their existing airport, to be funded by taxpayers. Comment on whether each of the following methods will produce a good random sample of the city's population:

a Surveying random travelers at the city's bus station.

b Picking random people from the city's phone book and surveying them.

c Surveying random employees of the existing airport.

d Surveying random patrons of an upscale mall.

3. (25pts) A repair shop counts how many vehicles come in each day for an oil change. The data is in the table below (it shows that 2 vehicles came on 5 days, 3 vehicles came on 13 days, etc.)

- a) Draw a histogram for the data.
- b) Find the mode number of daily oil changes.
- c) Find the median number of daily oil changes.
- d) Find the mean number of daily oil changes.
- e) Find the standard deviation.

| Oil changes | Frequency (days) |
|-------------|------------------|
| 2 | 5 |
| 3 | 13 |
| 4 | 17 |
| 5 | 11 |
| 6 | 3 |

4. (6pts) 205 lb Rodolfo is from Brazil, where the weight of men his age is normally distributed with mean 175 lbs and standard deviation 18 lbs. 188 lb Kiran is from India, where the weight of men his age is normally distributed with mean 164 lbs and standard deviation 13 lbs. Use z -scores to determine who is more overweight relative to the populations of their respective countries.

7. (17pts) When elevated to angle 45° , a gun can shoot a shell about 18 kilometers. Actually, tests with many firings of identical shells resulted in a normal distribution of distances with mean 18 km and standard deviation 0.75 km. Draw a picture showing which area you are computing as you answer:

- a) What percentage of shells fall farther than 19km away?
- b) What percentage of shells fall between 17.5 and 18.5 kilometers away?

Bonus. (10pts) Weights of watermelons from a certain field are normally distributed. If it is known that 25% of watermelons have weight under 9 lbs and 75% of watermelons have weight under 12.5 lbs find the mean weight of watermelons and the standard deviation.

(Hint: a part of this problem is the inverse of what we usually do: an area is given and we have to find the z -score. Once you have z -scores, the standard deviation is not far behind.)