All our formulas

No, you may not bring these to an exam

Rules for exponents:

 $a^{-n} = \frac{1}{a^n}$ $a^m \cdot a^n = a^{m+n}$ $\frac{a^m}{a^n} = a^{m-n}$ $(a^m)^n = a^{mn}$ $(ab)^m = a^m b^m$ $\left(\frac{a}{b}\right)^m = \frac{a^m}{b^m}$

Rules for roots:

$$\sqrt[n]{ab} = \sqrt[n]{a} \sqrt[n]{b}$$
$$\sqrt[n]{\frac{a}{b}} = \frac{\sqrt[n]{a}}{\sqrt[n]{b}}$$
$$\sqrt[n]{a^n} = |a|, \text{ for even } n$$
$$\sqrt[n]{a^n} = a, \text{ for odd } n$$
$$a^{\frac{m}{n}} = \sqrt[n]{a^m} = (\sqrt[n]{a})^m$$

Distance and midpoint

$$d = |a - b|$$

$$m = \frac{a + b}{2}$$

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

$$M = \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right)$$

Rules for logarithms:

$$\log_a a^x = x \qquad a^{\log_a x} = x$$
$$\log_b M = \frac{\log_a M}{\log_a b}$$
$$\log_a(MN) = \log_a M + \log_a N$$
$$\log_a \left(\frac{M}{N}\right) = \log_a M - \log_a N$$
$$\log_a M^p = p \cdot \log_a M$$

Algebraic expressions:

$(a+b)^2 = a^2 + 2ab + b^2$	square of a sum
$(a-b)^2 = a^2 - 2ab + b^2$	square of a difference
$a^2 - b^2 = (a - b)(a + b)$	difference of squares
$(a+b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$	cube of a sum
$(a-b)^3 = a^3 - 3a^2b + 3ab^2 - b^3$	cube of a difference
$a^3 - b^3 = (a - b)(a^2 + ab + b^2)$	difference of cubes
$a^3 + b^3 = (a+b)(a^2 - ab + b^2)$	sum of cubes

Circles and lines:

$(x-h)^2 + (y-k)^2 =$	r^2 circle with center (h, k) and radius r
$m = \frac{y_2 - y_1}{x_2 - x_1}$	slope of line through (x_1, y_1) and (x_2, y_2)
y = mx + b	line with slope m and y -intercept b
$y - y_1 = m(x - x_1)$	line with slope m through (x_1, y_1)

distance between real numbers a and b

midpoint of real numbers a and b

distance between points in the plane (x_1, y_1) and (x_2, y_2) midpoint of points in the plane (x_1, y_1) and (x_2, y_2)

Quadratic formula:

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Even, odd functions: If f(-x) = f(x), f is even If f(-x) = -f(x), f is odd

Compound interest:

$$A = P\left(1 + \frac{r}{n}\right)^{nt}$$