

1. (6pts) For the matrices A , B and C find the following expressions, if they are defined:

a) A^2C

b) BB^T

c) $2C - BA$

$$A = \begin{bmatrix} 2 & 1 \\ -1 & -1 \\ 0 & 3 \end{bmatrix}$$

$$B = \begin{bmatrix} 7 & 0 & 1 \\ -2 & 3 & 2 \end{bmatrix}$$

$$C = \begin{bmatrix} 2 & 1 \\ -1 & -1 \end{bmatrix}$$

2. (6pts) The matrix A is given below.

a) Find the inverse of A .

b) Use the inverse to effortlessly solve the system below.

$$A = \begin{bmatrix} 2 & 4 \\ 7 & -1 \end{bmatrix}$$

$$2x_1 + 4x_2 = 1$$

$$7x_1 - x_2 = 3$$

3. (4pts) Find the cosine of the angle between the vectors $\mathbf{a} = (1, -1, 3, 4)$ and $\mathbf{b} = (0, 4, 5, 2)$.

4. (9pts) A system of linear equations is given below.

a) Use the Gauss-Jordan method (that is, transform the augmented matrix to reduced row-echelon form) in order to solve the system.

b) Write the solution in vector form.

c) Describe the set of points in \mathbf{R}^4 that the solution set represents.

$$3x_1 + x_2 + 13x_4 = 11$$

$$-x_2 - x_3 - 6x_4 = -1$$

$$2x_1 + 2x_2 + x_3 + 17x_4 = 9$$

5. (5pts) Below is the augmented matrix of a system of linear equations. Determine the c 's for which the system has: a) one solution, b) infinitely many solutions, c) no solutions. (Note: no row operations are needed.)

$$A = \begin{bmatrix} 1 & 3 & 4 & 5 + c \\ 0 & 1 & -17 & 7 \\ 0 & 0 & c^2 - 4c & c - 4 \end{bmatrix}$$

6. (3pts) The matrix B was obtained by applying a row operation to matrix A . Find the elementary matrix E so that $EA = B$.

$$A = \begin{bmatrix} 3 & 7 & -7 \\ 1 & -6 & 4 \end{bmatrix} \quad B = \begin{bmatrix} 7 & -17 & 9 \\ 1 & -6 & 4 \end{bmatrix} \quad E =$$

7. (3pts) Find a 2×2 matrix B so that $B \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} 3a & 3b \\ 5a & 5b \end{bmatrix}$ for every 2×2 matrix.

8. (4pts) Suppose we have a system with 4 equations in 3 unknowns. Every equation represents a plane in \mathbf{R}^3 . Draw one example of a 4-plane arrangement for each of the following situations: a) the system has no solution b) the solution is a line in \mathbf{R}^3 .

9. (10pts) Are the following statements true or false? Justify your answer by giving a logical argument or a counterexample.

a) If \mathbf{u} is orthogonal to \mathbf{v} , then $\|\mathbf{u} + \mathbf{v}\|^2 = \|\mathbf{u} - \mathbf{v}\|^2$

b) If A is a 3×5 matrix with at least 2 non-zero entries, then the solution set of the linear system $A\mathbf{x} = \mathbf{0}$ always has at most 3 parameters.

c) If A is an $n \times n$ matrix and $A^{17} = I$, then A is invertible.

Bonus. (5pts) Use a linear system to show that the vector $(-2, 25, 11)$ is a linear combination of vectors $(2, 1, 3)$ and $(3, -5, 1)$ and find the coefficients that realize this linear combination.