

Calculus 2 — Exam 5
MAT 308, Fall 2021 — D. Ivanišić

Name: _____
Show all your work!

1. (12pts) Polar coordinates of two points are given.
- Sketch the points in the plane.
 - For each point, give two additional polar coordinates, one with a negative r , one with a negative θ .

$$\left(-1, \frac{3\pi}{4}\right) \qquad \left(3, \frac{7\pi}{5}\right)$$


2. (10pts) Convert (a picture may help):
- $\left(8, \frac{2\pi}{3}\right)$ from polar to rectangular coordinates
 - $(-2\sqrt{3}, 2)$ from rectangular to polar coordinates

3. (14pts) Find the equation of the tangent line to the parametric curve $x = \frac{\cos t}{t}$, $y = \frac{\sin t}{t}$ at the point where $t = \pi$.

4. (12pts) A particle moves along the path with parametric equations $x(t) = \sin t$, $y(t) = 3 + \cos^2 t$ for $0 \leq t \leq 2\pi$. Eliminate the parameter in order to sketch the path of motion and then describe the motion of the particle.

5. (6pts) Identify the curve given in polar coordinates by $r = 5 \sec \theta$ by converting the equation to cartesian coordinates.

6. (12pts) The graph of $r = f(\theta)$ is given in cartesian coordinates. Use its intervals of increase and decrease to help you sketch the polar curve $r = f(\theta)$. Indicate which piece of the cartesian graph corresponds to which piece of the polar graph.



7. (16pts) The parametric curve $x = 3 + t$, $y = 1 + 2t$, $-1 \leq t \leq 2$ is given.

- Use an integral to find the length of the curve.
- Eliminate the parameter in order to find out what the curve is.
- Use a college algebra method to find its length and compare your answer to a).

8. (18pts) A parametric curve is given by $x(t) = 2t^3 - 3t^2$, $y(t) = t^2 - 6t + 2$.

a) Find the points on the curve where the tangent line is horizontal or vertical.

b) Where does the curve go as $t \rightarrow \infty$ and $t \rightarrow -\infty$? (That is, find $\lim_{t \rightarrow \pm\infty} x(t)$, $\lim_{t \rightarrow \pm\infty} y(t)$.)

c) Plot the points from a) on a coordinate system and use them, along with information from b), or from plotting additional points, to get a graph of the curve. Recall that the curve moves in only one of general directions $\nearrow \nwarrow \swarrow \searrow$ between points from a).

Bonus. (10pts) Show that the length of a polar curve $r = f(\theta)$, $\alpha \leq \theta \leq \beta$ is given by the formula below. *Hint: to start, get a parametrization $(x(\theta), y(\theta))$ of the polar curve by using formulas for the polar \rightarrow cartesian conversion.*

$$l = \int_{\alpha}^{\beta} \sqrt{f(\theta)^2 + f'(\theta)^2} d\theta.$$